



Castelnuovo-Mumford Regularity of Toric Surfaces

Sean Grate

Auburn University

January 11, 2025

Get bored?

Check out my website (seangrate.com)

- Other math
 - Lefschetz properties
 - algebraic combinatorics
 - topological data analysis
 - software
- I'm on the job market looking for a postdoc position
 - CV
 - research statement
 - teaching and learning statement

Betti tables

Given an ideal $I \subseteq R = \mathbb{K}[x_0, \dots, x_n]$, a free resolution “approximates” R/I . The Betti table records the ranks of the summands appearing in a free resolution.

Definition

Let

$$0 \leftarrow R/I \leftarrow F_0 \leftarrow F_1 \leftarrow \dots F_n \leftarrow 0$$

be the minimal free resolution of R/I with $F_i = \bigoplus_j R(-j)^{\beta_{i,j}}$. The Betti table of F_\bullet is a table where the entry in the j -th row and i -th column is $\beta_{i,i+j}$.

Example

$R = \mathbb{K}[x, y, z]$ and $I = \langle x^2, y^2, z^3 \rangle$

$$0 \leftarrow R/I \leftarrow R \xleftarrow[\oplus]{R(-3)}^{(x^2 \ y^2 \ z^3)} R(-2)^2 \xleftarrow[\oplus]{R(-5)^2}^{\begin{pmatrix} -y^2 & -z^3 & 0 \\ x^2 & 0 & -z^3 \\ 0 & x^2 & y^2 \end{pmatrix}} R(-4) \xleftarrow[\oplus]{R(-5)^2}^{\begin{pmatrix} z^3 \\ -y^2 \\ x^2 \end{pmatrix}} R(-7) \leftarrow 0$$

	0	1	2	3
0	1	.	.	.
1	.	2	.	.
2	.	1	1	.
3	.	.	2	.
4	.	.	.	1

Example

$R = \mathbb{K}[x, y, z]$ and $I = \langle x^2, y^2, z^3 \rangle$

$$0 \leftarrow R/I \leftarrow R \leftarrow \begin{array}{c} R(-(1+1))^2 \\ \oplus \\ R(-(1+2)) \end{array} \leftarrow \begin{array}{c} R(-(2+2)) \\ \oplus \\ R(-(2+3))^2 \end{array} \leftarrow R(-(3+4)) \leftarrow 0$$

	0	1	2	3
0	1	.	.	.
1	.	2	.	.
2	.	1	1	.
3	.	.	2	.
4	.	.	.	1

Castelnuovo-Mumford regularity

Definition

Let $I \subseteq \mathbb{K}[x_0, \dots, x_n]$ be a homogeneous ideal, and consider the minimal free resolution

$$0 \leftarrow R/I \leftarrow F_0 \leftarrow F_1 \leftarrow \cdots \leftarrow F_{n+1} \leftarrow 0$$

of R/I , where $F_i \cong \bigoplus_j R(-i-j)^{\beta_{i,i+j}}$. The Castelnuovo-Mumford regularity (or simply regularity) is

$$\text{reg}(R/I) = \max_{i,j} \{j : \beta_{i,i+j} \neq 0\}.$$

The regularity is the index of the bottom row of the Betti table.

Castelnuovo-Mumford regularity

Definition

A coherent sheaf \mathcal{F} on \mathbb{P}^n is m -regular if

$$H^i(\mathcal{F}(m-i)) = 0$$

for all $i > 0$. The Castelnuovo-Mumford regularity (or simply regularity) is

$$\inf \{d : H^i(\mathcal{F}(d-i)) = 0 \text{ for all } i > 0\}.$$

Monomial curves

Definition

The monomial curve with exponents $a_1 \leq \dots \leq a_{n-1}$ in \mathbb{P}^n is the curve $C \subset \mathbb{P}^n$ of degree $d = a_n$ parameterized by

$$\varphi: \mathbb{P}^1 \rightarrow \mathbb{P}^n \quad \text{with} \quad (s, t) \mapsto (s^d, s^{d-a_1}t^{a_1}, \dots, s^{d-a_{n-1}}t^{a_{n-1}}, t^d).$$


Theorem (L'vovsky; 1996)

Let $A = (0, a_1, \dots, a_n)$ be a sequence of non-negative integers such that the g.c.d. of the a_j 's equals 1, and let C be the corresponding monomial curve. Then C is m -regular, where

$$m = \max_{1 \leq i < j \leq n} \{(a_i - a_{i-1}) + (a_j - a_{j-1})\},$$

i.e., m is the sum of the two largest gaps in the semigroup generated by A .

Example (twisted cubic)



$$A = \begin{pmatrix} 0 & 1 & 2 & 3 \end{pmatrix} \longrightarrow \begin{pmatrix} 0 & 1 & 2 & 3 \\ 3 & 2 & 1 & 0 \end{pmatrix}$$

$$\varphi(s, t) = (s^3, \underset{x_0}{s^2t}, \underset{x_1}{st^2}, \underset{x_2}{t^3})$$

$$I_C = \langle x_2^2 - x_1x_3, \quad x_1x_2 - x_0x_3, \quad x_1^2 - x_0x_2 \rangle$$

$$0 \leftarrow R/I_C \leftarrow R \leftarrow R(-2)^3 \leftarrow R(-3)^2 \leftarrow 0$$

$$\text{reg}(I_C) = 2 \leq (a_1 - a_0) + (a_2 - a_1) = 1 + 1 = 2$$

```
i1 : kk = ZZ/32749;

i2 : I = monomialCurveIdeal(kk[x_0..x_3], {1,2,3})

o2 = ideal (x2 - x1x3, x2x1 - x2x2, x2 - x0x3)
          2           2
          2   1 3   1 2   0 3   1   0 2

i3 : print betti res I
      0 1 2
total: 1 3 2
      0: 1 . .
      1: . 3 2
```

Example (sporadic)



$$A = \begin{pmatrix} 0 & 3 & 5 & 7 \end{pmatrix} \rightarrow \begin{pmatrix} 0 & 3 & 5 & 7 \\ 7 & 4 & 2 & 0 \end{pmatrix}$$

$$\varphi(s, t) = (s^7, \underset{x_0}{s^3 t^4}, \underset{x_1}{s^5 t^2}, \underset{x_2}{t^7})$$

$$I_C = \langle x_2^2 - x_1 x_3, \quad x_1^3 x_2 - x_0^2 x_3^2, \quad x_1^4 - x_0^2 x_2 x_3 \rangle$$

$$0 \leftarrow R/I_C \leftarrow R \leftarrow \frac{R(-2)}{\oplus R(-4)^2} \leftarrow R(-5)^2 \leftarrow 0$$

$$\text{reg}(I_C) = 4 \leq (a_1 - a_0) + (a_2 - a_1) = 3 + 2 = 5$$

```
i1: kk = ZZ/32749;
```

```
i2: I = monomialCurveIdeal(kk[x_0..x_3], {3,5,7})
```

```
o2 = ideal (x2 - x3x1, x3x2 - x2x3, x4 - x2x3x1)  
          2      3      2 2    4      2  
          1 3      1 2      0 3      1      0 2 3
```

```
i3 : print betti res I
```

```
0 1 2
```

```
total: 1 3 2
```



```
0: 1 . .
```

```
1: . 1 .
```

```
2: . . .
```

```
3: . 2 2
```

Toric surfaces




Goal

Extend L'vovsky's result to toric surfaces, i.e., find a combinatorial bound on the regularity of toric surfaces.

- *Look at toric surfaces which are defined by an incomplete linear series.*
- *Include all points on the boundary of a polygon and exclude all its interior points.*
- *These are usually not normal, but may be smooth.*

Setup



$$A = \begin{pmatrix} 0 & 0 & 3 \\ 0 & 3 & 0 \end{pmatrix}$$

↓ Conv(A) \ Int(A)

$$\begin{pmatrix} 0 & 1 & 2 & \cdots & 0 & 0 & 0 \\ 0 & 0 & 0 & \cdots & 3 & 2 & 1 \end{pmatrix}$$



↓ homogenize

$$\tilde{A} = \begin{pmatrix} 0 & 1 & 2 & \cdots & 0 & 0 & 0 \\ 0 & 0 & 0 & \cdots & 3 & 2 & 1 \\ 3 & 2 & 1 & \cdots & 0 & 1 & 2 \end{pmatrix}$$

Hollow triangle

Definition

Suppose $A = \begin{pmatrix} 0 & k & 0 \\ 0 & 0 & k \end{pmatrix}$. The hollow triangle of length k is $\Delta^k := \tilde{A}$.



Hollow triangle data

		0	1	2	3				
2	total:	1	6	8	3				
	0:	1	.	.	.				
	1:	.	6	8	3				
		0	1	2	3	4	5	6	7
3	total:	1	17	53	91	108	83	37	9
	0:	1
	1:	.	17	43	36	8	.	.	.
	2:	.	.	10	55	100	83	37	9

Hollow triangle data

```
+-----+
| |          0   1   2   3   4   5   6   7   8   9   10  11
| 4 | total: 1  33  153  525 1356 2178 2205 1486 675  201  36   3
| |          0: 1   .   .   .   .   .   .   .   .   .   .   .
| |          1: .   33  123  144   30   .   .   .   .   .   .   .
| |          2: .   .   30  381 1326 2178 2205 1486 675  201  36   3
| |
+-----+
```

```
+-----+
| |          0   1   2   3   4   5   6   7   8   9
| 5 | total: 1  54  389 2028 7845 18957 30393 34672 29106 18162
| |          0: 1   .   .   .   .   .   .   .   .   .
| |          1: .   54  266  462  174   15   .   .   .   .
| |          2: .   .   123 1566 7671 18942 30393 34672 29106 18162
| |
+-----+
```

Results

Lemma

For all $d \geq 2$, $(R/I_{\Delta^k})_d = (\overline{R/I_{\Delta^k}})_d$.

Theorem

For all $k \geq 2$, $\text{reg}(\Delta^k) = 2$.

Results

Lemma

For all $d \geq 2$, $(R/I_{\square^k})_d = (\overline{R/I_{\square^k}})_d$.

Theorem

For all $k \geq 2$, $\text{reg}(\square^k) = 2$.

Proof sketch of theorem

- Use the short exact sequence of sheaves

$$0 \rightarrow \mathcal{I}_{\square^k}(d) \rightarrow \mathcal{O}_{\mathbb{P}^{4k-1}}(d) \rightarrow \mathcal{O}_{\square^k}(d) \rightarrow 0$$


to eventually get a short exact sequence

$$0 \rightarrow R/I_{\square^k} \rightarrow \overline{R/I_{\square^k}} \rightarrow N \rightarrow 0.$$

- By the lemma, N is generated is only generated by degree 1 monomials.
- $\text{reg}(R/I_{\square^k}) \leq \max(\text{reg}(\overline{R/I_{\square^k}}), N) = \text{reg}(\overline{R/I_{\square^k}}) = 2$.

Proof sketch of lemma


- Showing $(R/I_{\square^k})_d = (\overline{R/I_{\square^k}})_d$ for $d \geq 2$ amounts to a computation with the lattice points of $\overline{\square^k}$.
- We are done if for any $p_1, p_2 \in \overline{\square^k}$, we can write $p_1 + p_2 = q_1 + q_2$ with $q_1, q_2 \in \square^k$.



$$p_1 + p_2 = \begin{pmatrix} 1 \\ 2 \\ 5 \end{pmatrix} + \begin{pmatrix} 2 \\ 2 \\ 4 \end{pmatrix} = \begin{pmatrix} 3 \\ 4 \\ 9 \end{pmatrix}$$

Proof sketch of lemma


- Showing $(R/I_{\square^k})_d = (\overline{R/I_{\square^k}})_d$ for $d \geq 2$ amounts to a computation with the lattice points of $\overline{\square^k}$.
- We are done if for any $p_1, p_2 \in \overline{\square^k}$, we can write $p_1 + p_2 = q_1 + q_2$ with $q_1, q_2 \in \square^k$.



$$\begin{aligned} p_1 + p_2 &= \begin{pmatrix} 1 \\ 2 \\ 5 \end{pmatrix} + \begin{pmatrix} 2 \\ 2 \\ 4 \end{pmatrix} = \begin{pmatrix} 3 \\ 4 \\ 9 \end{pmatrix} \\ &= \begin{pmatrix} 0 \\ 2 \\ 6 \end{pmatrix} + \begin{pmatrix} 3 \\ 2 \\ 3 \end{pmatrix} \end{aligned}$$

Proof sketch of lemma

- Showing $(R/I_{\square^k})_d = (\overline{R/I_{\square^k}})_d$ for $d \geq 2$ amounts to a computation with the lattice points of $\overline{\square^k}$.
- We are done if for any $p_1, p_2 \in \overline{\square^k}$, we can write $p_1 + p_2 = q_1 + q_2$ with $q_1, q_2 \in \square^k$.




$$\begin{aligned} p_1 + p_2 &= \begin{pmatrix} 1 \\ 2 \\ 5 \end{pmatrix} + \begin{pmatrix} 2 \\ 2 \\ 4 \end{pmatrix} = \begin{pmatrix} 3 \\ 4 \\ 9 \end{pmatrix} \\ &= \begin{pmatrix} 0 \\ 2 \\ 6 \end{pmatrix} + \begin{pmatrix} 3 \\ 2 \\ 3 \end{pmatrix} \\ &= \begin{pmatrix} 0 \\ 0 \\ 8 \end{pmatrix} + \begin{pmatrix} 3 \\ 4 \\ 1 \end{pmatrix} \end{aligned}$$


“Bad Boy”

In general, the regularity can be arbitrarily large by using

$$A = \tilde{A} = \begin{pmatrix} 0 & d & d-1 & d \\ 0 & d-1 & d & d \end{pmatrix}.$$




Smooth is not enough



	0	1	2	3	4	5	...
total:	1	54	385	1462	3608	6456	...
0:	1
1:	.	52	280	730	1128	1050	...
2:	.	.	81	600	2040	4416	...
3:	.	2	24	132	440	990	...

Smooth is not enough



	0	1	2	3	4	5	...
total:	1	74	633	2883	8593	18953	...
0:	1
1:	.	73	486	1627	3388	4620	...
2:	.	1	144	1218	4983	13541	...
3:	.	.	3	38	222	792	...

Thank you!