


Incidence toric ideals and three-point functions

Barbara Betti · Sean Grate · Thiago Holleben · Flavio Salizzoni

SIAM TX-LA Sectional Meeting



The Crew




Barbara Betti



Sean Grate



Thiago Holleben



Flavio Salizzoni

Three-point functions


Massless momenta $p_1, \dots, p_n \in \mathbb{C}^d$, with
Mandelstam invariants $p_{ij} = p_i \cdot p_j$
(Lorentzian, $p_{ii} = 0$).

Three-point functions, all scaling dimensions 2:

$$C_{ijk} = \frac{1}{p_{ij} p_{jk} p_{ik}}.$$

Question


Which algebraic relations hold among the C_{ijk} ?



$$c_{ijk} := C_{ijk}^{-1} = p_{ij} p_{jk} p_{ik}$$

The answer depends on the dimension

Let $\tilde{I}_{n,d} \subseteq \mathbb{C}[c_{ijk}]$ be the ideal of relations when the momenta live in $\mathbb{C}^d$.



Toric for $d \geq n$: Rajan–Sverrisdóttir–Sturmfels (2024). Gram-matrix minors for $d < n$: El Maazouz–Sturmfels–Sverrisdóttir (2024).


Incidence toric ideals

Definition

$$\varphi: \mathbb{K}[c_\kappa : \kappa \in \binom{[n]}{k}] \longrightarrow \mathbb{K}[x_\tau : \tau \in \binom{[n]}{t}], \quad c_\kappa \longmapsto \prod_{\tau \subset \kappa, |\tau|=t} x_\tau$$

$I_{n,k,t} := \ker \varphi$, with exponent matrix $A_{n,k,t}$ and polytope


$P_{n,k,t} := \text{conv}(\text{columns of } A_{n,k,t})$.



$$k = 3, t = 2: \tilde{I}_{n,n} = I_{n,3,2}$$

- $t = k - 1$: a signless boundary map $\rightsquigarrow$ topology
- $t = 1$: the hypersimplex $\Delta(k, n)$
- homogeneous: every column has $\binom{k}{t}$ ones

Running example: $n = 6, k = 3, t = 2$



antipodal pairs $\{1, 2\}, \{3, 4\}, \{5, 6\}$

30

minimal generators

5

codimension

162


degree

15 *octahedral quartics*, one for each octahedron on $[6]$:

$$c_{135} c_{146} c_{236} c_{245} - c_{136} c_{145} c_{235} c_{246}$$

The two sides are the blue and orange facets. Plus 15 sextics.

A Lefschetz map in disguise



$n = 4$: monomials of $A \leftrightarrow$ subsets. Blue:
 $\times L: A_2 \rightarrow A_3$.

$$A = \mathbb{K}[x_1, \dots, x_n] / (x_1^2, \dots, x_n^2)$$

and $L = x_1 + \dots + x_n$. Multiplying by L adds one element in every possible way, so

$$\times L^{k-t}: A_t \rightarrow A_k \text{ is } (k-t)! A_{n,k,t}^\top$$

(cf. Holleben 2025).

Theorem (Stanley 1980)

In characteristic 0, monomial complete intersections have the strong Lefschetz property.

Proposition

$A_{n,k,t}$ has full rank:

$$\text{rank } A_{n,k,t} = \min\left\{\binom{n}{t}, \binom{n}{k}\right\}.$$

Proof. $\times L^{k-t} = (k-t)! A_{n,k,t}^T$ has full rank by Stanley (1980).

Consequences. $I_{n,k,t} \neq 0$ iff $\binom{n}{t} < \binom{n}{k}$, and then $\text{ht } I_{n,k,t} = \binom{n}{k} - \binom{n}{t}$,
 $\dim V(I_{n,k,t}) = \binom{n}{t} - 1$.

Example. $I_{6,3,2}$ has height $20 - 15 = 5$.

Characteristic p and the polytope

Theorem (Jonsson Kling 2026)

In characteristic p , the map $\times L^{k-t}: A_t \rightarrow A_k$ has full rank if and only if $p > \min(k, n - t)$.

Proposition

Every prime $p \leq \min(k, n - k + 1)$ divides the normalized Euclidean volume of $P_{n,k,k-1}$.

Corollary

The product of all primes $p \leq k$ divides $\text{Vol}(P_{2k-1,k,k-1})$, so the volume is not a polynomial in n, k, t .

Caveat. Euclidean lattice, not the lattice of $A_{n,k,t}$: this is not $\deg V(I_{n,k,t})$, which remains open.

Binomials are null designs

Definition

$f: \binom{[n]}{k} \rightarrow \mathbb{Z}$ is a *null t -design* if $\sum_{F \supseteq X} f(F) = 0$ for every t -subset X .

$$\ker_{\mathbb{Z}} A_{n,k,t} \cong \{\text{null } t\text{-designs}\}, \quad f \mapsto c^{f+} - c^{f-}$$

Theorem (Deza–Frankl–Singhi 1983)

A nonzero null t -design takes at least 2^t positive values.

So every binomial in $I_{n,k,t}$ has degree at least 2^t .


Pods: the degree bound is sharp

**Theorem (Graver–Jurkat 1973;
Graham–Li–Li 1980)**

Null t -designs are spanned by the *pods*

$$\prod_{m=0}^t (x_{i_{2m+1}} - x_{i_{2m+2}}) \prod_{m=2t+3}^{k+t+1} x_{i_m}.$$

A pod has exactly 2^t positive terms, so the lowest degree in $I_{n,k,t}$ is exactly 2^t .



$(x_1 - x_2)(x_3 - x_4)(x_5 - x_6)$: terms are cube vertices, sign is parity — the octahedral quartic.

Theorem

$P_{n,k,t}$ is $(2^t - 1)$ -neighborly,
and not 2^t -neighborly.

Proof. A non-face F yields $v \in \ker A_{n,k,t}$ with $\text{supp}(v_+) \subseteq F$, so $|F| \geq 2^t$ by Deza–Frankl–Singhi (1983). The positive half of a pod is a non-face of size exactly 2^t .

Example. $P_{6,3,2}$ has 20 vertices in dimension 14: every 3 vertices span a face, but the 4 blue triangles of an octahedron do not.

Octahedra generate


Pods are the two-row Specht polynomials (Morita–Wachi–Watanabe 2009), i.e. orientations of hyperoctahedra.

Theorem (Spink 2020)

$\ker_{\mathbb{Z}} A_{n,k,t}$ is generated by orientations of hyperoctahedra.

So the *octahedral ideal* $J_{n,k,t}$ they generate satisfies

$$I_{n,k,t} = J_{n,k,t} : \left(\prod_{\kappa} c_{\kappa} \right)^{\infty}.$$



Do the extra components matter?

Question


Do the components of $V(J_{n,k,t})$ in the coordinate hyperplanes have lower dimension, so that $\deg V(I_{n,k,t}) = \deg V(J_{n,k,t})$?

The smallest case $(6, 3, 2)$ in Macaulay2:

	codim	degree
$V(J_{6,3,2})$	5	162
$V(I_{6,3,2})$	5	162

Yes [here](#): no extra component of the top dimension.

Balanced complexes



Colour classes $\{1, 2\}$, $\{3, 4\}$, $\{5, 6\}$; facets 2-coloured blue/orange.

For $t = k - 1$, $A_{n,k,k-1}$ is the top boundary map of the simplex with the signs forgotten.

Definition (Stanley 1979)

Δ is *balanced* if its vertices can be k -coloured so that every facet sees each colour once.

Lemma (Klee–Novik 2016)

A balanced normal pseudomanifold without boundary, of dimension ≥ 2 , is orientable iff its facet–ridge graph is bipartite.

Theorem


Balanced orientable pseudomanifolds
give Graver basis elements.

Let $\Delta = \langle \sigma_1, \dots, \sigma_s \rangle$ be a balanced, orientable, normal $(k-1)$ -pseudomanifold without boundary on $[n]$, $k \geq 3$, with orientation ε . Then

$$\prod_{\varepsilon_i=1} c_{\sigma_i} - \prod_{\varepsilon_j=-1} c_{\sigma_j}$$


lies in the Graver basis of $I_{n,k,k-1}$.

Proof idea. Order vertices by colour: signs of ∂ depend only on the deleted colour, so $D\partial$ is a submatrix of $A_{n,k,k-1}$, with kernel $\tilde{H}_{k-1}(\Delta; \mathbb{Z}) = \mathbb{Z}\varepsilon$.



The octahedron gives the quartics.

New Graver elements from cross-flips



This disk replaces the triangle 136 of the octahedron.

Cross-flips (Izmestiev–Klee–Novik 2017) preserve balancedness and topology.

Octahedron $\rightsquigarrow$ balanced 2-sphere with 9 vertices and 14 triangles $\rightsquigarrow$

$C_{146} C_{236} C_{135} C_{245} C_{678} C_{179} C_{389}$

— $C_{789} C_{167} C_{368} C_{139} C_{246} C_{145} C_{235}$

a degree-7 Graver element of $I_{9,3,2}$.

Open questions

Degree

Compute $\deg V(I_{n,k,t})$.

Octahedra vs. toric

Is $\deg V(I) = \deg V(J)$ in general? Yes for $(6, 3, 2)$.

Non-toric

Find bases for $\tilde{I}_{n,d}$ with $d < n - 1$.

Lefschetz

rank, dimension, volume
arithmetic

Null designs

degree bound, sharp
neighborliness

Topology

Graver elements from
balanced spheres